

УДК 551.465

© V. G. Gnevyshev<sup>1</sup>, V. S. Travkin<sup>2</sup>, T. V. Belonenko<sup>2\*</sup>, 2023

© Translation from Russian: E. V. Rykova, 2023

<sup>1</sup>Shirshov Institute of Oceanology, Russian Academy of Sciences, 36 Nakhimovsky Prosp., Moscow, 117997, Russia

<sup>2</sup>St. Petersburg State University, 7–9 Universitetskaya Emb., St. Petersburg, 199034, Russia

\*btvlibab@yandex.ru

## TOPOGRAPHIC FACTOR AND LIMIT TRANSITIONS IN THE EQUATIONS FOR SUB-INERTIAL WAVES

Received 29.09.2022, Revised 08.11.2022, Accepted 08.12.2022

### Abstract

In this paper, sub-inertial waves propagating on the Kuril shelf and the oceanic trench are considered. Against the background of a historical review of the beginning of the study of topographic waves and the appearance of relevant terms, a description of the features of wave propagation and the derivation of the main dispersion equations are given. We show that all variants of the topographic solutions presented in the article are basically based on the same dispersion relation: this is the dispersion relation for Rossby topographic waves. Two separate classes of localized solutions have been constructed: one is for the shelf, and the second, in fact, is also for the shelf, but which is commonly called trench waves. We demonstrate that the transverse wave number for trench waves is not independent, as for shelf waves, but is a function of the longitudinal wave number. In other words, Rossby topographic waves are two-dimensional waves, while shelf waves are quasi-one-dimensional solutions. The analytical novelty of the work consists of the fact that we can make crosslinking of trench and shelf waves. This fact was not presented in previous articles on this topic.

**Keywords:** shelf, ocean trench, topographic waves, shelf waves, trench waves, crosslinking solutions

© В. Г. Гневывшев<sup>1</sup>, В. С. Травкин<sup>2</sup>, Т. В. Белоненко<sup>2\*</sup>, 2023

© Перевод с русского: Е. В. Рыкова, 2023

<sup>1</sup>Институт океанологии им. П.П. Ширшова РАН, 117997, Россия, Москва, Нахимовский проспект, 36.

<sup>2</sup>Санкт-Петербургский государственный университет, 199034, Санкт-Петербург, Россия,

Университетская наб., 7–9, г.

\*btvlibab@yandex.ru

## ТОПОГРАФИЧЕСКИЙ ФАКТОР И ПРЕДЕЛЬНЫЕ ПЕРЕХОДЫ В УРАВНЕНИЯХ ДЛЯ СУБИНЕРЦИОННЫХ ВОЛН

Статья поступила в редакцию 29.09.2022, после доработки 08.11.2022, принята в печать 08.12.2022

### Аннотация

Рассматриваются топографические субинерционные волны, распространяющиеся на шельфе и океаническом желобе. На фоне обзора истории исследования топографических волн и появления соответствующих терминов авторы дают описание особенностей распространения этих волн и вывод основных дисперсионных уравнений. Показано, что все варианты представленных в статье решений в основе своей базируются на одном и том же дисперсионном соотношении — это дисперсионное соотношение для топографических волн Россби. Построены два класса локализованных решений: одно для шельфовых волн, второе, фактически, тоже шельфовое, но его принято называть желобовыми волнами. Показано, что для желобовых волн поперечное волновое число не является независимым, как для шельфовых волн, а является функцией от продольного волнового числа. Другими словами, топографические волны Россби — это всегда двумерные волны, в то время как шельфовые волны представлены квазиодномерными решениями. Аналитическая новизна работы состоит в том, что в ней удалось произвести сшивки желобовых и шельфовых волн, которые ранее отсутствовали в работах по данной тематике.

**Ключевые слова:** шельф, океанский желоб, волны топографические, шельфовые, желобовые, сшивки решений

Ссылка для цитирования: Гневывшев В.Г., Травкин В.С., Белоненко Т.В. Топографический фактор и предельные переходы в уравнениях для субинерционных волн // Фундаментальная и прикладная гидрофизика. 2023. Т. 16, № 1. С. 8–23. doi:10.48612/fpg/92rg-6t7h-m4a2

For citation: Gnevyshev V.G., Travkin V.S., Belonenko T.V. Topographic Factor and Limit Transitions in the Equations for Sub-inertial Waves. *Fundamental and Applied Hydrophysics*. 2023, 16, 1, 8–23. doi:10.48612/fpg/92rg-6t7h-m4a2

## 1. Introduction

In recent years, much research has appeared that studies various classes of sub-inertial waves in the ocean in areas characterized by significant changes in bottom topography (see, for example, [1–5]). For these ocean areas, wave energy capture is observed by huge topographic changes. Abrupt depth changes in the open ocean lead to the appearance of “topographic” waves, which are similar in nature to the shelf waves formed in the zone of the shelf-continental slope [6]. The comparative analysis of the contribution of various factors to the dispersion equations shows that the topographic factor is dominant [2–5, 7, 8]. Shelf waves are the most important type of sub-inertial oscillations, which plays an important role in the variability of the ocean level in the synoptic frequency range [6, 9].

In the fields of deep-sea trenches, another type of sub-inertial oscillation is formed, the propagating of which along the ocean side of the trench forms a system of trench waves. The term “trench waves” was first used by Mysak et al. [10] to study sub-inertial waves propagating in ocean trenches (Kuril, Japanese, Japanese-Kuril, Chilean, Peruvian). In the mathematical model of trench waves, two-dimensional topography is assumed to be uniform along the slope of the trench and has terminal scales of transverse variability.

To describe linear trench waves, a mathematical model is used, which is based on the line Schrödinger equation with additional boundary conditions that lead to quantization of the transverse wave number. The solution is constructed in a trigonometric form (sine), localized across the topographic feature, and a monochromatic plane wave traveling along the topographic feature — the slope of the trench. Note that the solution of the type “Kelvin double wave”, for which two exponents are used in the topographic profile, is not considered in the trench wave model [10].

If the two-dimensional derivative of the shelf depth is constant, then mathematically the problem allows us to analyze the boundary problem in general form and formulate certain theorems of the spectral problem [6]. Solutions to this problem are called “shelf waves” [11].

If the two-dimensional derivative of the shelf depth changes a sign when moving away from the shore (ocean trench), then the mathematical analysis is significantly complicated. For trench-type topography, particular classes of piecewise smooth analytical solutions are constructed, which are conventionally called “trench waves”. A feature of many ocean trenches is their close position to the coastline, thereby combining the shelf and continental slope areas and adjoining the inside of the trench (Figure 1). In the future, we will call the inner part of the trench, consisting of the shelf and the continental slope, reaching the maximum depth of the trench, “trench shelf”. The oceanic side of the trench has less variability in depth and, as a rule, greater steepness [12]. Therefore, in the construction of the mathematical model of trench waves, the entire physical semiinfinite space is conditionally divided into three regions: 1) the trench shelf where the depth of the ocean increases; 2) the oceanic part of the trench, where the depth of the ocean decreases and the derivative of the function describing the depth changes the sign; 3) an open ocean, where the depth of the ocean is constant and finite.


Since only sub-inertial oscillations are considered trench wave model [10], the Earth’s rotation rate is considered constant (there is no  $\beta$ -effect) and there are no shear flows, then the solution is always decaying exponentially in the open part of the ocean, and this solution is sometimes called “exponential tail”.

We will consider wave movements on the  $f$ -plane, i. e.

$$f = \text{const.} \quad (1.1)$$

The consistency of the Earth’s rotation rate destroys waves with frequencies less than the frequency of rotation of the Earth itself. Thus, in the open part of the ocean, sub-inertial wave oscillations are suppressed. This fact was first noted by Longuet-Higgins [13, 14].

In addition, the trench wave model [10] examines for sub-inertial oscillations with spatial scales of hundreds of kilometers, but does not take into account the  $\beta$ -effect in the open part of the ocean, i. e. a strictly closed wave system is built, and Rossby waves falling on the shelf from the open part of the ocean are actually excluded from the problem. The latter is the main shortage of this model. According to the theory [10, 11, 15], trench waves can be represented in the form of two particular solutions: 1) shelf waves: the solution is localized on the shelf part of the trench, while in the direction from the shelf towards the open ocean this solution can be represented



**Fig. 1.** Profile of the oceanic trench: 1 — trench shelf (shelf + continental slope), 2 — oceanic part of the trench, 3 — open ocean

as oscillation-attenuation-attenuation; 2) trench waves: the solution is localized on the oceanic part of the trench, this solution can be imagined as attenuation—oscillation—attenuation. A fundamentally important point is that in at least one part of the profile in the trench wave model (recall that there are three parts of the profile — trench shelf, trench, open ocean) there must be an oscillatory solution, moreover, there must be a sine in the solution, in the other two must be a damped exponential function, and in the open part of the ocean an exponential tail [10]. In this case, as shown in the paper [10], the shelf part of the solution is asymptotically stitched with the Buchwald and Adams shelf wave [16], if the model eliminates the oceanic trench part of the topography and turns its width to zero. However, acting similarly for the chute part of the solution and rushing the width of the already shelf part of the topography to zero, the authors [10] could not sew the chute solution with the inner shelf wave of Buchwald and Adams. They simply did not analyze this fact, citing little interest in this class of solutions. As a result, the question: of whether trench waves are sewn or not sewn [10] with the decision of Buchwald and Adams [16] remaining open. In this work, we will eliminate this drawback and show that the trench solution [10] is sewn together with the inner shelf wave of Buchwald and Adams. To do this, it is necessary not to zero the width of the shelf part of the topography (the trench shelf), placing the solid wall in the center of the trench, as it was done [10], but, on the contrary, to push it infinitely far. As a result, the analysis of trench waves becomes logical and complete.

Thus, we formulate the purpose of the study as follows: we will show that the solution of the “trench waves” is asymptotically sewn together with the topographic and internal topographic waves of Buchwald and Adams. The specificity consists in the method of the limit transition, namely: to sew the oceanic part of the trench solution of Mysak et al. [10] with the inner shelf wave of Buchwald and Adams, you need to direct the trench part of the solution to zero, or, conversely, direct the width of the trench to infinity while simultaneously striving for the steepness of the shelf to zero. And to crosslink the trench wave with the internal topographic wave of Buchwald and Adams, you also need to endlessly move the border, expanding the shelf area and turning zero shelf steepness.

This work is constructed as follows. In the second section, we give the basic equations and give the dispersion relations for the Rossby topographic waves. Next, we see into question from simple to complex: first in the third section, we give the Buchwald and Adams solution, which subsequently becomes the asymptotics of the more complex solution of Mysak et al. Then, in the fourth section, we give this solution [10] for trench waves, simultaneously eliminating many misprints in their work, and we sew successfully the solution [10] asymptotically with the Buchwald and Adams solution. As a result, we prove that the Buchwald and Adams solution is the asymptotics of the solution of Mysak et al. [10].

The central idea of the article, which is discussed in the next three sections (5–7), is the statement that all variants of the topographic solutions presented in the article are based on the same dispersion relation — this is the dispersion relation for topographic Rossby waves. We show that this approach greatly simplifies the perception of the different topographic models discussed in the work.

## 2. Formulation of the problem. Basic equations

The theoretical basis for describing trench waves is linearized barotropic equations of shallow water in the rigid-lid approximation taking into account the topography [6, 10]:

$$u_t - fv + \rho^{-1}p_x = 0, \quad (2.1)$$

$$v_t + fu + \rho^{-1}p_y = 0, \quad (2.2)$$

$$(Hu)_x + (Hv)_y = 0. \quad (2.3)$$

Following the work [10], we assume the geostrophic approximation, thereby filtering out the high-frequency continuous spectrum of Poincaré waves, as well as edge waves:

$$u = -\Psi_y/H, \quad v = -\Psi_x/H. \quad (2.4)$$

The following designations are used:  $u$  and  $v$  are the velocity components in the  $x$  and  $y$  directions,  $\Psi$  — current function,  $H$  is the depth,  $p$  is the pressure,  $\rho$  is the water density, and  $f$  is the Coriolis parameter. The classical right coordinate system is adopted, axes  $x$  and  $y$  are directed across the trench (away from the coast ( $x = -L_1$ )) and along the trench, respectively.

Substituting (2.4) in (2.1) and (2.2) (equation (2.3) is performed automatically) and considering solutions of wave nature along the topographic feature (trench slope)

$$\Psi = \Psi(x) \exp i(ky - \omega t), \quad (2.5)$$

we get the following linear homogeneous one-dimensional equation:

$$\left(\frac{\Psi_x}{H}\right)_x + \left[\frac{f k}{\omega} \left(\frac{1}{H}\right)_x - \frac{k^2}{H}\right] \Psi = 0, \quad (2.6)$$

where  $k$  — is the longitudinal component of the wave number, which is strictly real and, according to [10], strictly positive,  $\omega$  — is the frequency of the wave, which is also real and can take both positive and negative values. Accordingly, depending on the sign, different directions of the phase velocity are obtained. Note that, unlike this approach, in the monograph [6], on the contrary, the  $\omega$  is always positive, but the wave number  $k$  can take both positive and negative values. Therefore, different directions of wave propagation — northeast or southwest — are achieved by a different sign of wave number.

The cross-linking conditions at  $x = 0$  (the center of the trench is the point of maximum depth on the profile) и  $x = L_2$  (the boundary of the trough on the side of the open ocean)

$$[\Psi] = 0, x = 0, x = L_2, \quad (2.7)$$

$$\left[\frac{\Psi_x + (f k / \omega) \Psi}{H}\right] = 0, \quad x = 0, \quad x = L_2. \quad (2.8)$$

Condition (2.8) for a topography model with a continuous function takes the form

$$[\Psi_x] = 0, x = 0, x = L_2. \quad (2.9)$$

Also, in the shelf area, the impermeability boundary condition is set at the point  $x = -L_1$  [10]:

$$[\Psi] = 0, x = -L_1 \quad (2.10)$$

and decaying condition at infinity:

$$\Psi \rightarrow 0, x \rightarrow \infty. \quad (2.11)$$

### *Dispersion relation for the Rossby topographic wave on the exponential profile of the topography*

The classical substitution  $\Psi = H^{1/2}$  in equation (2.6) can remove the first derivative [3, 9] and bring the above equation to the equation with the classical harmonic oscillator potential. The most successful and easy-to-analyze oceanic trough model is one with an exponential topography profile under the analytical approach and one with a stepped profile for numerical counting. There are other special cases of parameterization of the topography when it is possible to obtain non-reflective solutions [17, 18], and solutions in terms of degenerate hypergeometric functions [19].

For the classical topographic Rossby wave on the exponential profile of topography  $H = H_0 \exp(-x/L)$ ,  $L$  is the shelf width, a solution is sought in the form of  $\Psi = H^{1/2} \exp[i(k_1 x + k_2 y - \omega t)]$ , where  $k_1$  and  $k_2$  are the transverse (across the trench) and longitudinal wavenumbers. Then the dispersion relation has the form [3, 9]:

$$\omega = \frac{-(\beta_0 + f/L)k_2}{k_1^2 + k_2^2 + 1/4L^2}, \quad (2.12)$$

where  $\beta_0$  is the  $\beta$ -plane parameter.

The main idea of this work is that all varieties of subinertial topographic waves, including shelf and trench waves, follow precisely from this dispersion relation. And therefore, we propose to call the relation (2.12) the general dispersion relation on the exponential profile of topography (see also [4, 5]).

Historically, the first to construct an analytical model of subinertial topographic waves were Buchwald and Adams [16]. In their work, they analyzed two qualitatively different models of topographic waves: shelf waves and internal shelf waves. The principal difference between an internal shelf wave and a conventional shelf wave is the presence of an infinitely extended ocean of finite depth. This additional degree of freedom leads to the appearance of a certain feature in the behavior of the 0-mode, the analysis of which becomes extremely sensitive in the long-wave limit. The latter can be explained by the fact that the first mode either passes into a Kelvin double wave, or both waves have close dispersion curves in the long wave range. Looking ahead, we note that Buchwald and Adams saw certain difficulties in the limiting transition and considered its different options.

At the same time, Longuet-Higgins [13] conducts an extremely laborious analysis using higher-order decompositions to achieve the physical correct behavior of the frequency (the tendency of the frequency to zero when striving for zero, and not to the wavenumber constant). Other authors, in particular, [6, 10, 11] take a different path using numerical counting, and show smooth dispersion curves, including for the first mode. This approach does not mean that the analysis applied is incorrect, but rather indicates that different options are possible in the long-wave limit, and the result depends on what exactly the author is trying to find, given the scheme by which the boundary problem is numerically considered.

We move on to the description of the first and simplest model of topographic Rossby waves.

### 3. Buchwald and Adams shelf waves

The topography in the model [16] is taken in the following exponential form (see Fig. 2):

$$\begin{aligned} H_1 &= H_0 \exp(2bx), \quad 0 < x \leq 1, \\ H_2 &= H_0 \exp(2b), \quad x > 1. \end{aligned} \quad (3.1)$$

We are looking for a solution in the following form

$$\begin{aligned} \Psi_1 &= A \sin m x \exp b(x-1), \quad 0 < x < 1, \\ \Psi_2 &= A \sin m \exp[-k(x-1)], \quad x \geq 1, \end{aligned} \quad (3.2)$$

where we believe  $k > 0$ . Then the dispersion relation (2.12) will take the form

$$\omega = \frac{-2f b k}{k^2 + m^2 + b^2}. \quad (3.3)$$

The wave amplitude in this case decays exponentially at the same rate on both sides of the break. Further, using cross-linking derivatives of functions, we obtain a condition for a transverse wavenumber:

$$\tan m = -\frac{m}{k+b}. \quad (3.4)$$

Further analysis will consider the sewing together of shelf and trench waves.

#### *Buchwald and Adam's internal shelf waves*

The trench wave model [10] assumes that the shelf width is finite and that the shelf near the shore extends to a finite depth. However, if we assume that the left boundary of the model has finite depth and is moved to infinity, then the boundary condition (2.10) is replaced by the attenuation criteria (2.11). This type of topography is called “rapid underwater slope” or simply “underwater slope” ([9], paragraph 24, page 311). In the model, the approach of the “rigid lid” is taken, while considering waves with lengths less than the barotropic Rossby radius. The phase associated with each of its eigenfunctions propagates so that shallow water remains on the right (in the northern half-globe). It can also be shown that these waves have dispersion. As shelf width approaches zero, higher modes are reduced to steady currents.

Following [16], we assume the following exponential topography model (Fig. 3):

$$\begin{aligned} H_1 &= h_1, \quad x \leq 0, \\ H_2 &= h_1 \exp 2bx, \quad 0 \leq x \leq L_2, \\ H_3 &= h_1 \exp 2bL_2, \quad L_2 \leq x < \infty. \end{aligned} \quad (3.5)$$

A solution is sought as follows:

$$\begin{aligned} \Psi_1 &= A_1 \exp(|k|x), \quad x \leq 0, \\ \Psi_2 &= (A_2 \sin mx + A_3 \cos mx) \exp(bx), \quad 0 \leq x \leq L_2, \\ \Psi_3 &= A_4 \exp(-|k|(x-L_2)), \quad L_2 \leq x < \infty. \end{aligned} \quad (3.6)$$




Fig. 2. Topography profile in Buchwald and Adam's model [16]



Again, using the dispersion relation for Rossby topographic waves, we obtain

$$\omega_n = \frac{-2f b k}{k^2 + l_n^2(k) + b^2}, \quad n = 1, 2, \dots \quad (3.7)$$

From the conditions of crosslinks, we get the following condition for the definition  $l_n(k)$ :

$$\tan L_2 l_n = -\frac{2l_n |k|}{l_n^2(k) + b^2 - k^2}, \quad (3.8)$$

where  $n$  — mode number.

The analysis performed in the study [16] shows that the internal shelf waves practically coincide with the “ordinary” shelf waves, except for the 0-mode. The 0-mode of internal shelf waves has a certain specificity: its internal transverse wave number  $l$  at small longitudinal wave numbers  $k$  becomes virtual  $l = i l_0$ . Accordingly, equation (3.8) takes the form:

$$\tanh L_2 l_0 = \frac{2l_0 |k|}{b^2 - l_0^2 - k^2}. \quad (3.9)$$

At the same time, the 0-mode frequency tends not to zero, as it happens in all other modes of topographic Rossby waves, but to the Earth’s rotation frequency [16], which looks somewhat strange. Looking ahead, we note that Longuet-Higgins [13, 14] received a different result, and below we will show it. When divergence is taken into account, the frequency, as in many physical models, tends to zero, and the result is classical long waves. In the limit transition, when the width of the topographic feature tends to zero:  $L_2 \rightarrow 0$ ,  $b \rightarrow 0$ ,  $bL_2 = 1/2 \ln(H_1/H_3)$  this 0-mode, according to [16], passes into a divergence-free double Kelvin wave. For this mode, the decision over the topographic feature is no longer sinusoidal, but exponential. The term “Kelvin double wave” belongs to Longuet-Higgins [14].

However, and this is fundamentally important: the Kelvin double wave appears in the problem of internal shelf waves precisely as the limit transition of the 0-mode when moving to the type of topography in the form of a step and has nothing to do with the trench waves of Mysak et al. Note also that a characteristic scale for the 0-mode appears in the problem:  $k_0 = (\sqrt{1 + b^2 L_2^2} - 1) / L_2$ . This is the scale of the solution change in the 0-mode when the sinusoidal solution over the trench goes into a double exponential, i. e. Kelvin double wave. Below you will see that a close scale will be discussed in the next section. For the Kuril trench, for example, the value  $bL_2 = 0.2332$ , (see Table 1 [10]); wavelength calculations give a wave number  $k_0 = 5 \times 10^{-4} \text{ km}^{-1}$ , which corresponds to a wavelength of approximately 13,000 km. This is too large a value, and the appearance of instability is purely theoretical.

#### 4. Mysak trench waves

In trench waves of Mysak et al. [10], the trench solution is highly localized above the topographic feature only from the shelf side. The value of the proper function on the shore has the order  $10^{-3}$  of the value above the topography [6]. Therefore, for it, replacing the left leakage boundary condition with attenuation physically should not have a significant effect. For a trench wave, the presence of an infinitely extended open ocean of finite depth plays a more important role.

This model considers a more complex configuration, which essentially consists of two types of waves. It combines the shelf waves and the internal shelf waves of Buchwald and Adams [16]. Next, following the work [10], we assume the following exponential model of the trench:

$$\begin{aligned} H_1 &= H_0 \exp 2\alpha x, & -L_1 \leq x \leq 0, \\ H_2 &= H_0 \exp(-2\beta x), & 0 \leq x \leq L_2, \\ H_3 &= H_0 \exp(-2\beta L_2), & L_2 \leq x < \infty. \end{aligned} \quad (4.1)$$




Fig. 3. Buchwald and Adam's topography model [16]

We have for the Kuril trench (see [10], Table 1):  $H_0 = 8.25$  km,  $L_1 = 185$  km,  $L_2 = 55$  km,  $\alpha = 0.95 \times 10^{-2}$  km $^{-1}$ ,  $\beta = 4.24 \times 10^{-3}$  km $^{-1}$ ,  $a = \alpha L_1 = 1.75$ ,  $b = \beta L_1 = 0.784$ ,  $r = L_2 / L_1 = 0.297$ .

The topography profile is continuous, however, the derivative of the topography profile suffers a break. Therefore, a solution for the current function is sought on the class of piecewise smooth functions using the crosslinking conditions at  $x = 0$  and  $x = L_2$ . The result obtained in work [10] can be formulated as follows: a solution satisfies automatically the condition (2.8) in the form

$$\begin{aligned}\Psi_1 &= A^{1/2} \exp(\alpha x) \quad -L_1 \leq x \leq 0, \\ \Psi_2 &= (B \sin kx + A \sin mL_1 \cos kx) \exp(-\beta x), \quad 0 \leq x \leq L_2, \\ \Psi_3 &= (B \sin(l L_2) + A \sin mL_1 \cos(l L_2)) \exp(-\beta L_2 - k(x - L_2)), \quad L_2 \leq x < \infty.\end{aligned}\quad (4.2)$$

Note that in the article [10] in the last formula there is a typo, which is corrected here. Further, due to the attenuation of the decision in the open part of the ocean, in formula (3.8) the wavenumber  $k$  is taken positive  $k > 0$ , and the frequency  $\omega$  can take both positive and negative values (in the monograph [6]  $\omega > 0$ , but  $k$  can be both positive and negative).

In (4.2) the parameters  $m$  and  $l$  are determined by the formulas:

$$m = \left( -\frac{2\alpha f k}{\omega} - \alpha^2 - k^2 \right)^{1/2}, \quad (4.3)$$

$$l = \left( \frac{2\beta f k}{\omega} - \beta^2 - k^2 \right)^{1/2}. \quad (4.4)$$

Substituting (4.2) in (2.8), we get a system of two homogeneous equations for  $A$  and  $B$ . The matrix of C coefficients has the form

$$C = \begin{pmatrix} \sin(mL_1) [(k - \beta) \cos(l L_2) - l \sin(l L_2)] & [(k - \beta) \sin(l L_2) + l \cos(l L_2)] \\ \sin(mL_1) \{ \alpha + \beta \} + m \cos(mL_1) & -l \end{pmatrix}. \quad (4.5)$$

Equating the determinants of the matrix to zero, [10] obtained the following equation:

$$(\alpha + \beta + m \cot mL_1) [l + (k - \beta) \tan l L_2] + l (k - \beta - l \tan l L_2) = 0. \quad (4.6)$$

Note that [10], as well as subsequently [6], call this equation a dispersion relation. Note that in formula (4.6) we corrected another typo in the last term of work [10].

Next, when moving from (4.5) to (4.6), authors [10] make an implicit assumption:  $\sin(mL_1) \neq 0$ ,  $\cos(lL_2) \neq 0$ . Thus, constructed localized solutions in work [10] exist, but do not exhaust all possible scenarios.

The numerical analysis of the relation (4.6) performed [10] is as follows: the complete spectral problem allows two partial approximate solutions. The first solution is shelf waves; the solution is oscillatory in the area of the shelf and the inside of the trench and decays in the rest of the physical space. The second solution is trench waves; the solution oscillates only in the area of the marine part of the trench, fading on the shelf and in the open ocean. The authors argue that these particular solutions (shelf waves and trench waves) can be obtained as limit transitions from a common trench model; they are localized in separate parts of the trench and are solutions to the common ocean trench model with very good accuracy.


The description of these private solutions is below.

## 5. Asymptotic of the trench waves on the shelf. Crosslinking with shelf wave

The main types of bottom relief, which determine the types of trapped waves, are shown in Figure 4. In all cases, waves propagate so that in the northern hemisphere (in the southern — on the contrary) shallow water (shore) remains on the right side relative to the direction of movement.

Let's start our consideration with a solution for the littoral zone. Suppose that on the shelf the solution is oscillating. Then the wavenumber  $m$ , given by the relation (4.3) must be positive. For this purpose,  $\omega < 0$  the condition must be met

$$\frac{-2\alpha k}{\alpha^2 + k^2} < \frac{\omega}{f} < 0. \quad (5.1)$$



**Fig. 4.** Types of captured waves (direction shown by arrows) depending on the shape of the bottom relief: *a* — trench waves; *b* — shelf waves; *c* — Kelvin waves; *d* — internal shelf waves; *e* — Kelvin double waves

Since the  $y$ -axis is directed north, we obtain from the expression for phase in (2.5) in the longitudinal direction of the trench, we receive  $\frac{\omega}{k} < 0$ . Therefore, the shelf wave phase propagates in a negative direction, i. e. to the southwest, along the shore on the right (northern hemisphere). In this case, from the relation (4.4), it is automatically obtained that the wave number  $l$  in the trench part will be purely imaginary, therefore, for the trench, the solution is damped.

The first qualitative solution for the shelf is oscillation-attenuation-attenuation (direction from the shore towards the open ocean). Authors [10] crosslinked the solution as follows: from (4.6) at the limit transition  $L_2 \rightarrow 0$  ( $\beta$  and  $l$  are limited) we get the equation

$$(k + \alpha) \sin m L_1 + m \cos m L_1 = 0 \quad (5.2)$$

or

$$\tan m L_1 = -\frac{m}{k + \alpha}, \quad m = m_n(k), \quad n = 1, 2, \dots \quad (5.3)$$

The ratio for shelf wave amplitudes  $B/A \rightarrow 0$ . Such a method of sewing, in which the width of the trench part tends to zero, gives the correct result due to the fact that on the right there is a semi-infinite ocean of finite depth. However, an attempt to do the same symmetrically for the shelf part, rushing the shelf width to zero, leads to an unexpected result with a singularity. Mysak et al. [10] did not comment on this singularity and left the question open. Therefore, for the symmetry of the limit transitions of the shelf and trench waves, we make an alternative limit transition at  $L_2 \rightarrow \infty$ ,  $\beta \rightarrow 0$ . At the same time, it turns out  $l \rightarrow ik$ , and taking advantage



of the ratio  $\tan i / L_2 = i \tanh l / L_2$ , we as a result again get the ratio (5.3). This result is extremely stable at  $L_2 \rightarrow \infty$ ,  $\beta \rightarrow 0$ , does not depend on the value of the quantity  $\beta L_2$ , which can be either infinitely large or infinitely small. Graphical analysis of equation (5.3) is performed in the work [16], and here we will not give it.

From the side of analytics, the main idea of this part of our work is to consider trench waves as some sub-species of classical Rossby topographic waves with a classical dispersion relation. We also believe that there is no great need to introduce new classes of waves with transcendental dispersion relations. Developing this direction, we offer an alternative interpretation to approaches [10] and [6]. We believe that equations (4.3) and (4.4) are more correctly called dispersion relations for topographic waves, and we consider the ratio (4.6) and its special cases as an additional condition that determines the transverse component of the wave number.

Thus, we get some subclasses of classical topographic Rossby waves with a dispersion relation in which there is a feature in finding the transverse component of the wave number. In fact, we propose to return to the interpretation that was proposed in the first original work on this topic i. e. [16].

### ***Shelf waves in an alternative presentation***

Let's give a simpler interpretation of the shelf part of the solution [10] — the shelf wave in terms of topographic Rossby waves. Recall that we are talking only about sub-inertial fluctuations.

Consider formula (4.1) in relation to the shelf part of the topography. Assume  $1/L = -2\alpha$ ,  $k_1 = m$ ,  $k_2 = k$ , and substitute it in (2.12). Finally, we obtain the relation (4.3) for shelf waves in the classical form of topographic Rossby waves in the form:

$$\omega = \frac{-2\alpha f k}{k^2 + m^2 + \alpha^2}. \quad (5.4)$$

It is this ratio that is more correctly called the dispersion relation for shelf waves. Further, it is necessary to impose the condition that the shelf is not infinite but has a finite width  $L_1$ . It is geometrically clear that then a certain condition will appear that will allow us to sew the sine on the shelf side of the trench with an exponent on the ocean part of the trench for each fixed value of the wavenumber  $k$ , where  $0 < k < \infty$  (according to [10],  $k$  is considered positive). To do this, you need to find the value of  $m$ , by solving equation (5.3), and then get a functional connection  $m = m(k)$ . There will be a number of such solutions  $m = m_n(k)$ ,  $n = 1, 2, \dots$ . Next, we substitute the values found in (5.4) and plot (we apply one point on the dispersion curve). Then we take a new value of the wavenumber  $k$  and repeat the procedure. Therefore, the dispersion relation for shelf waves should be recorded as

$$\omega = \frac{-2\alpha f k}{k^2 + m_n^2(k) + \alpha^2}, \quad n = 1, 2, \dots \quad (5.5)$$

Based on the result of the work [16], we will give an assessment for  $m_n(k)$ :

$$\frac{(n-1/2)\pi}{L_1} < m_n(0) \leq m_n(k) \leq \frac{n\pi}{L_1}. \quad (5.6)$$

Functional connection  $m = m(k)$  extremely low changes the dispersion curve of topographic Rossby waves. Note that [10] and [16] did not consider one special case, which should also be implemented. The condition  $\sin(mL_1) = 0$  is equivalent to setting an additional solid wall at  $x = 0$ :  $\Psi_1(0) = \Psi_2(0) = 0$ . For the shelf zone, the wave number  $m$  ceases to depend on the longitudinal component of the wave vector  $k$ , so we immediately find a discrete spectrum in an explicit form:

$$\sin(mL_1) = 0, \quad m = \frac{n\pi}{L_1}, \quad n = 1, 2, \dots \quad (5.7)$$

The wave number  $l$  in the trench part is again purely imaginary, i. e. attenuation outside the shelf occurs. The dispersion relation (4.3) takes the form:

$$\omega = -\frac{2\alpha f k}{k^2 + m^2 + \alpha^2} = -\frac{2\alpha f k}{k^2 + (n\pi / L_1)^2 + \alpha^2}. \quad (5.8)$$

This relation is qualitatively similar to (5.5), but it is simpler in the sense that it does not contain an unknown dependence  $m = m(k)$  and thus does not require any numerical counting. This dispersion relation is of a particular nature, but in practice, it can be useful due to its simple form and quite understandable geometric meaning:  $n$  is the number of sine half-periods that fit into the shelf zone.

Next, following [16], we find the long-wave asymptotic  $k \rightarrow \infty$  of the dispersion relation (5.5):

$$\omega \approx \frac{-2\alpha f k}{m_n^2(0) + \alpha^2}, \quad n = 1, 2, \dots \quad (5.9)$$

The dependence on the wave number turned out to be linear, as it should be in the long-wave limit. Then the phase velocity is equal to the group velocity and equal to

$$c = c_{gr} \approx \frac{-2\alpha f}{m_n^2(0) + \alpha^2}, \quad n = 1, 2, \dots \quad (5.10)$$

Let's move on to dimensionless variables, taking the scale of length  $L_1$  is the width of the shelf part of the trench: parameter  $\alpha^* = \alpha L_1$ , the longitudinal wave number  $k^* = k L_1$ , the transverse wave number  $m^* = m L_1$ . Next, for ease of recording, omit the asterisks of the variables. Then the estimate of, for example, the first transverse wavenumber (the first mode) has the form:

$$\frac{\pi}{2} < m_1 \leq \pi, \quad (5.11)$$

The second mode

$$\frac{3\pi}{2} < m_2 \leq 2\pi. \quad (5.12)$$

For the Kuril profile, the dispersion curves of shelf and trench waves are presented in Figure 5.

## 6. Features of trench waves

Let's pay attention to the next point related to terminology. The monograph [6] (p. 102) states the following: [10] considered "Kelvin double waves propagating along the ocean side of the trench to the northeast". However, the term "Kelvin double waves" is not found at all in [10], and the authors call the solutions obtained for the "trench waves". We also adhere to the term "trench waves" [20, 21]. The term "Kelvin double waves" was introduced by Longuet-Higgins [13, 14], as the limit transition of the 0-mode on a continuous topographic feature to a topographic profile of the "step" type when the width of the topographic feature




Fig. 5. Dispersion diagram calculated using the formula (5.8) for the Kuril region: shelf (left) and trench (right) waves

tends to zero. The maximum transition made below in our work shows the validity of the use of this term. Indeed, let the solution be oscillating in the trench, i. e. the wavenumber  $l$  is real. Then it follows from (4.4) that the relation:

$$0 < \frac{\omega}{f} < \frac{2\beta k}{\beta^2 + k^2}. \quad (6.1)$$

In this case, the wavenumber  $m$  automatically becomes purely imaginary  $m = iM$ , where  $M = \left( \frac{2\alpha f k}{\omega} + \alpha^2 + k^2 \right)^{1/2}$  (a positive root sign is taken). The dispersion relation in the limit  $L_1 \rightarrow 0$  ( $\alpha$  — limited) has the form:

$$(k - \beta) \sin(l L_2) + l \cos(l L_2) = 0 \quad (6.2)$$

or (see [10])

$$\tan(l L_2) = -\frac{l}{k - \beta}. \quad (6.3)$$

Also, at  $k = \beta$  [10]  $l = 0$  — a trivial solution is believed. However, from (6.2) it is clear that there is also a non-trivial solution  $\cos(l L_2) = 0$ . Then the amplitude ratio for the trench waves is inverted and becomes  $A/B = 0$ .

### ***Trench waves in alternative presentment***

Let's make the following transformations: in the second formula (4.1) we assume  $1/L = 2\beta$ ,  $k_1 = l$ ,  $k_2 = k$ , and then substitute it in (2.12). Then, for the exponential topography profile, consider the dispersion relation for topographic waves

$$\omega = \frac{2\beta f k}{k^2 + l^2 + \beta^2}. \quad (6.4)$$

But (6.4) is also a dispersion relation for trench waves. Next, for each fixed value of the wavenumber  $k > 0$ , you need to find the value of  $l$ , by solving equation (6.2). Thus, we get a dependence  $l = l(k)$ . There will be a countable set of such solutions  $l = l_n(k)$ ,  $n = 1, 2, \dots$ , so the dispersion relation for trench waves is more correct to write in the form

$$\omega_n = \frac{2\beta f k}{k^2 + l_n^2(k) + \beta^2}, \quad n = 1, 2, \dots \quad (6.5)$$

Further, for numerical counting, we will again move on to dimensionless variables, only taking as the length scale the width of the oceanic part of the trench  $L_2$ :  $l^* = l L_2$ ,  $k^* = k L_2$ ,  $\beta^* = \beta L_2$ . For the Kuril trench, we get  $\beta^* = 0,2332$ . As before, we omit asterisks for variables. Then the equation (6.3) for finding the dependence in dimensionless form for the Kuril trench takes the form

$$\tan l = -\frac{l}{k - 0.2332}. \quad (6.6)$$

This wavenumber  $l$  has already been discussed above, and since it gives a very large wavelength, it is not of practical interest.

For trench waves in the Kuril trench, an important role has the presence of an open ocean of constant depth on the outside of the trench. According to the results of the work [15], which is also confirmed by the results [16], the open ocean has little effect on short trench waves, while long waves most likely become unstable. Perhaps it is this instability that gives rise to the vortex formations observed in the Kuril trench [22–27].

The limit transition  $L_1 \rightarrow 0$  for the trench wave, made by the authors [10], physically means the transfer of the solid wall from the shelf to the deepest part of the trench. It is not clear how to avoid the singularity for the resulting asymptotics. It also looks somewhat strange from a physical point of view, since a limit transition is made for the shelf wave  $L_2 \rightarrow \infty$ , and the ocean (right) boundary of the trench is moved to infinity. At the same time, for the shelf part of the solution, the expected result is obtained, and the shelf asymptotic enters the shelf wave.

Now let's consider an alternative transition for trench asymptotics. Instead  $L_1 \rightarrow 0$  let's assume, on the contrary:  $L_1 \rightarrow \infty$ , we move the shelf (left) border of the trench to infinity. In this case, we also add a new condition  $\alpha \rightarrow 0$ , the steepness of the topography tends to zero. Then the component of the wavenumber  $m$  becomes purely imaginary. For certainty, assume  $m = ik$ ,  $k > 0$ . Then, taking into account  $\cot ik = -i \coth k$ , the relation (4.6) for determining the transverse component of the wavenumber  $l$  takes the form:

$$\tan l L_2 = \frac{2l k}{l^2 + \beta^2 - k^2}. \quad (6.7)$$

And this is a familiar equation for internal shelf waves, obtained [16]. Thus, we get a completely understandable physical result: in the same way as the shelf asymptotic [10] gives the shelf wave [16], the trench, the oceanic part of the solution of the trench goes to the internal shelf wave [16]. It is unclear why Mysak et al. [10] did not show this moment; it may not have worked out because they shifted the hard wall to the center of the trench.

## 7. Kelvin double wave

The classical interpretation of Kelvin double waves is based on a stepped model of topography [9]:

$$\begin{aligned} H &= H_1, & x < 0, \\ H &= H_2, & x > 0. \end{aligned} \quad (7.1)$$

In this case, equation (2.6) will take the form

$$\Psi_{xx} - k^2 \Psi = 0. \quad (7.2)$$

The solution satisfying the attenuation at infinity has a simple form:

$$\begin{aligned} \Psi &= A \exp(|k| x), & x < 0, \\ \Psi &= A \exp(-|k| x), & x > 0. \end{aligned} \quad (7.3)$$

The wave amplitude in this case decays exponentially at the same rate on both sides of the break.

Using the sewing condition, the following dispersion relation is obtained:

$$\frac{\omega}{f} = \frac{H_2 - H_1}{H_2 + H_1} \operatorname{sgn}|k|, \quad (7.4)$$

where  $H_2 > H_1$ , and  $\operatorname{sgn}$  is the sign change function. Then for a profile of the “step” type, it turns out that for any wave number, there is only one solution i. e. one subinertial mode. Note that the group velocity of these waves is always zero.

Although the term “Kelvin double wave” belongs to Longuet-Higgins [13, 14], the dispersion relation itself (7.4) seems to have been known even earlier and is also given in the work [16]. Later, Longuet-Higgins [13] considered the problem with continuous topography profiles and, using the limit transition from a continuous profile to a “step” profile, derived the second dispersion equation for Kelvin double waves. At the same time, it turned out that for a continuous profile, there is an enumerable setup of its eigenfunctions, but at the limit transition to the “step” profile, only one lower mode is insensitive to the profile shape in depth. Higher modes depend on the details of the profile and at the limit transition tend to stationary currents, on the basis of which Longuet-Higgins [13] concluded that the lowest mode is most likely to be observed in the real ocean.

A fundamentally important point is that in the next article, Longuet-Higgins [14] abandoned the dispersion relation (7.4). The reason is understandable: indeed, from a physical point of view, it is wrong if the phase velocities of long waves take infinitely large values. Longuet-Higgins [14], using more accurate (up to the second order) asymptotic decompositions, obtained that for the lower Rossby long wave mode, it is quite justified that the frequency tends to zero, and not to a constant when considering wave divergence. Essentially, for the barotropic mode, he added Rossby's barotropic radius to the denominator  $\operatorname{Россби} f^2/gH$ . However, in the case where the depth profile  $h(x)$  in the transition zone is exponential, this is already in contradiction with the result of Buchwald and Adams [16]. According to Longuet-Higgins [14], this contradiction is due to the fact that he took into account the horizontal divergence of wave motion. Note that these facts are not reflected in the works [6, 10, 11], although the dispersion curves constructed on the basis of numerical evaluation in the monograph

[6] correspond to the results of the Longuet-Higgins article [14]. One of these results is that the frequency tends to zero in the long wave limit. It should be noted that the limit transition procedure itself is not entirely unambiguous. The curvature of the dispersion curve for the 0-mode of a Kelvin double wave is extremely complex in the vicinity of zero (for long waves). Buchwald and Adams also note this [16]. Our estimates for the Kamchatka trench give too small wave number values, i. e. the corresponding wavelengths are most likely purely theoretical in nature since they actually go beyond the scale limits of the  $f$ -plane model.

In the second article of Longuet-Higgins [14], the author corrected his previous result, but it is not said in the monograph [9]. The main result is as follows: for all modes, including the lower mode (this is important), provided that wave divergence is taken into account, the group velocity of topographic Rossby waves and their phase velocity coincide in direction for long waves (with small wave numbers) and have opposite directions for short waves.

## 8. Conclusion

The following main conclusions can be drawn from the exponential topographic model of the trench. The phase of linear topographic low-frequency waves always propagates, leaving shallow water (or shore) on the right. Further, two separate classes of localized solutions can be built: one offshore, the second, in fact, also offshore, but which is usually called trench waves (in some sources this decision is mistakenly called Kelvin double waves). On the dispersion curves of both types of waves, there are both extremes (zero group velocity) and inflection points (group velocity extremum). This indicates that energy (vortices) can propagate along the shelf, both, for example, for the Kuril shelf, to the southwest (longer waves), and the northeast (shorter waves).

However, for the ocean part of the trench (for trench waves), this is not the case. Longer waves have a predominantly northeast direction of energy propagation, and shorter waves have a southwest direction. Trench waves are an order of magnitude slower (periods longer and speeds smaller) than shelf waves. This is qualitatively explained by the direct proportionality of the phase velocity of the slope of the exponential topography profile [4, 5].

Analytical models and numerical counting show that trench waves fade rather quickly toward the shelf and do not actually have a significant effect on the shelf part of the solution. Trench waves cannot be recorded while on shore, and when analyzing purely offshore solutions, it is possible not to take into account possible oscillation solutions localized on the trench.

Conversely, the shelf wave decays extremely slowly towards the trench (at least for the first mod numbers), which is extremely negative for trench localized solutions. We cannot say unequivocally that the shelf does not affect trench waves. Obviously, the effect of the shelf on the solutions localized at the trench can be manifested in a non-linear case.

The possibility of propagating topographic-type low-frequency Rossby waves captured by the Kurilo-Kamchatka trench to the pole was theoretically predicted in the studies [10, 26], which used the term “Kelvin double waves”. These quasi-geostrophic waves arise due to the persistence of potential vorticity and recovery mechanisms provided by the Coriolis force and the change in ocean depth.

The analytical novelty of this work is that we managed to cross-link the trench waves [10] and the shelf waves [16], physically justified and quite expected. To do this, we proposed to infinitely increase (expand) the corresponding areas of the shelf or trench while simultaneously striving to the steepness of the topography to zero.

## Funding

The publication was made with the financial support of the St Petersburg State University Grant No. 93016972, RSF Grant No. 22-27-00004, and State assignment theme No. 0128-2021-0003.

## References

1. Drivdal M., Weber J.E.H., Debernard J.B. Dispersion relation for continental shelf waves when the shallow shelf part has an arbitrary width: Application to the shelf west of Norway. *Journal of Physical Oceanography*. 2016, 46, 2, 537–549. doi:10.1175/jpo-d-15-0023.1
2. Gnevyshev V.G., Frolova A.V., Koldunov A.V., Belonenko T.V. Topographic effect for Rossby waves on a zonal shear flow. *Fundamental and Applied Hydrophysics*. 2021, 14, 1, 4–14. doi:10.7868/S2073667321010019



3. Pedlosky J. Geophysical fluid dynamics. Berlin, Springer, 1979. 624 p.
4. Gnevyshev V.V., Frolova A.V., Belonenko T.V. Topographic effect for Rossby waves on non-zonal shear flow. *Water Resources*. 2022, 49, 2, 240–248. doi:10.1134/S0097807822020063
5. Gnevyshev V.G., Frolova A.V., Kubryakov A.A., Sobko Yu.V., Belonenko T.V. Interaction between Rossby waves and a jet flow: Basic equations and verification for the Antarctic circumpolar current. *Izvestiya, Atmospheric and Oceanic Physics*. 2019, 55(5), 412–422. doi:10.1134/S0001433819050074
6. Efimov V.V., Kulikov E.A., Rabinovich A.B., Fine I.V. Waves in the ocean boundary regions. Leningrad, 1985, 250 p. (in Russian).
7. Belonenko T.V., Frolova A.V. Antarctic circumpolar current as a waveguide for Rossby waves and mesoscale eddies. *Sovremennye Problemy Distsionnogo Zondirovaniya Zemli iz Kosmosa*. 2019, 16, 1, 181–190. doi:10.21046/2070-7401-2019-16-1-181-190 (in Russian).
8. Belonenko T., Frolova A., Gnevyshev V. Detection of waveguide for Rossby waves using satellite altimetry in the Antarctic Circumpolar Current. *International Journal of Remote Sensing*. 2020, 41:16, 6232–6247, doi:10.1080/01431161.2020.1752955
9. LeBlond P., Mysak L.A. Waves in the Ocean. Elsevier Scientific Publishing Company, 1977. 602 p.
10. Mysak L.A., LeBlond P.H., Emery W.J. Trench Waves. *Journal of Physical Oceanography*. 1979, 9(5), 1001–1013. doi:10.1175/1520-0485(1979)009<1001: TW>2.0.CO;2
11. Mysak L.A. Recent advances in shelf wave dynamics. *Reviews of Geophysics and Space Physics*. 1980, 18, 1, 211–241.
12. Sandalyuk N.V., Gnevyshev V.G., Belonenko T.V., Kochnev A.V. Application of the vortex layer problem to the Gulf Stream area. *Sovremennye Problemy Distsionnogo Zondirovaniya Zemli iz Kosmosa*. 2021, 18, 5, 242–251. doi:10.21046/2070-7401-2021-18-5-242-251 (in Russian).
13. Longuet-Higgins M.S. On the trapping of waves along a discontinuity of depth in a rotating ocean. *Journal of Fluid Mechanics*. 1968, 31, 3, 417–434.
14. Longuet-Higgins M.S. Double Kelvin waves with continuous depth profiles. *Journal of Fluid Mechanics*. 1968, 34(01), 49. doi:10.1017/s002211206800176x
15. Mysak L.A., Johnson E.R., Hsiem W.W. Baroclinic and barotropic instabilities of coastal currents. *Journal of Physical Oceanography*. 1981, 11(2), 209–230. doi:10.1175/1520-0485(1981)011<0209>2.0.CO;2
16. Buchwald V.T., Adams J.K. The propagation of continental shelf waves. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*. 1968. 305(1481), 235–250. doi:10.1098/rspa.1968.0115
17. Pelinovsky E.N. Hydrodynamics of tsunami waves. Nizhny Novgorod. IAP RAS. 1996, 156 p. (in Russian).
18. Didenkulova I., Pelinovsky E. On shallow water rogue wave formation in strongly inhomogeneous channels. *Journal of Physics A: Mathematical and Theoretical*. 2016, 49(19), 194001. doi:10.1088/1751-8113/49/19/194001
19. Ermakov A.M., Stepanyants Y.A. Transformation of long surface and tsunami-like waves in the ocean with a variable bathymetry. *Pure and Applied Geophysics*. 2019, 177(3), 1675–1693. doi:10.1007/s00024-019-02259-4
20. Rabinovich A.B., Thomson R.E. Evidence of diurnal shelf waves in satellite-tracked drifter trajectories off the Kuril Islands. *Journal of Physical Oceanography*. 2001, 31, 2650–2668. doi:10.1175/1520-0485(2001)031<2650:EODSWI>2.0.CO;2
21. Rabinovich A.B., Thomson R.E., Bograd S.J. Drifter observations of anticyclonic eddies near Bussol' Strait, the Kuril Islands. *Journal of Oceanography*. 2002, 58, 661–671. doi:10.1023/A:1022890222516
22. Bulatov N.V., Lobanov V.B. Investigation of mesoscale eddies to the east of the Kuril Islands on the base of meteorological satellites data. *Issledovanie Zemli iz Kosmosa*. 1983, 3, 40–47 (in Russian).
23. Rogachev K.A. Rapid thermohaline transition in the Pacific western subarctic and Oyashio fresh core eddies. *Journal of Geophysical Research: Oceans*. 2000, 105, 8513–8526. doi:10.1029/1999JC900330
24. Kaneko H., Itoh S., Kouketsu S., Okunishi T., Hosoda S., Suga T. Evolution and modulation of a poleward propagating anticyclonic eddy along the Japan and Kuril-Kamchatka trenches. *Journal of Geophysical Research: Oceans*. 2015, 120, 4418–4440. doi:10.1002/2014JC010693
25. Prants S.V., Lobanov V.B., Budyansky M.V., Uleysky M. Yu. Lagrangian analysis of formation, structure, evolution and splitting of anticyclonic Kuril eddies. *Deep Sea Research, Part I*. 2016, 109, 61–75. doi:10.1016/j.dsr.2016.01.003
26. Efimov V.V., Rabinovich A.B. On resonant tidal currents and their connection with continental shelf waves in the northwestern Pacific Ocean. *Izvestiya Akademii Nauk SSSR. Fizika Atmosfery i Okeana*. 1980, 16, 10, 1091–1101 (in Russian).
27. Prants S.V. Trench eddies in the Northwest Pacific: An overview. *Izvestiya, Atmospheric and Oceanic Physics*. 2021, 57, 4, 341–353. doi:10.1134/S0001433821040216

## Литература

1. Drivdal M., Weber J.E.H., Debernard J.B. Dispersion relation for continental shelf waves when the shallow shelf part has an arbitrary width: Application to the shelf west of Norway // *Journal of Physical Oceanography*. 2016. Vol. 46, Iss. 2. P. 537–549. doi:10.1175/jpo-d-15-0023.1
2. Гневyshev В.Г., Фролова А.В., Колдунов А.В., Белоненко Т.В. Топографический эффект для волн Россби на зональном сдвиговом потоке // *Фундаментальная и прикладная гидрофизика*. 2021. Т. 14, № 1. С. 4–14. doi:10.7868/S2073667321010019
3. Pedlosky J. *Geophysical fluid dynamics*. Berlin: Springer, 1979. 624 p.
4. Gnevyshev V.V., Frolova A.V., Belonenko T.V. Topographic effect for Rossby waves on non-zonal shear flow // *Water Resources*. 2022. Vol. 49, N 2. P. 240–248. doi:10.1134/S0097807822020063
5. Гневyshev В.Г., Фролова А.В., Кубряков А.А., Собко Ю.В., Белоненко Т.В. Взаимодействие волн Россби со струйным потоком: основные уравнения и их верификация для Антарктического циркумполярного течения // *Известия РАН. Физика атмосферы и океана*. 2019. Т. 55, № 5. С. 39–50. doi:10.31857/S0002-351555539-50
6. Ефимов В.В., Куликов Е.А., Рабинович А.Б., Файн И.В. Волны в пограничных областях океана. Л.: Гидрометеиздат., 1985. 250 с.
7. Белоненко Т.В., Фролова А.В. Антарктическое циркумполярное течение как волновод для волн Россби и мезомасштабных вихрей // *Современные проблемы дистанционного зондирования Земли из космоса*. 2019. Т. 16, № 1. С. 181–190. doi:10.21046/2070-7401-2019-16-1-181-190
8. Belonenko T., Frolova A., Gnevyshev V. Detection of waveguide for Rossby waves using satellite altimetry in the Antarctic Circumpolar Current // *International Journal of Remote Sensing*. 2020. 41:16, P. 6232–6247, doi:10.1080/01431161.2020.1752955
9. Ле Блон П., Майсек Л. Волны в океане, в 2-х частях / Пер. с англ. М.: Мир, 1981. 846 с.
10. Mysak L.A., Leblond P.H., Emery W.J. Trench Waves // *Journal of Physical Oceanography*. 1979. Vol. 9, N 5. P. 1001–1013. doi:10.1175/1520-0485(1979)009<1001: TW>2.0.CO;2
11. Mysak L.A. Recent advances in shelf wave dynamics // *Reviews of Geophysics and Space Physics*. 1980. Vol. 18, N 1. P. 211–241.
12. Сандалюк Н.В., Гневyshev В.Г., Белоненко Т.В., Кочнев А.В. Приложение задачи о вихревом слое для района течения Гольфстрим // *Современные проблемы дистанционного зондирования Земли из космоса*. 2021. Т. 18, № 5. С. 242–251. doi:10.21046/2070-7401-2021-18-5-242-251
13. Longuet-Higgins M.S. On the trapping of waves along a discontinuity of depth in a rotating ocean // *Journal of Fluid Mechanics*. 1968. Vol. 31, pt. 3. P. 417–434.
14. Longuet-Higgins M.S. Double Kelvin waves with continuous depth profiles // *Journal of Fluid Mechanics*. 1968. Vol. 34, N 01. P. 49. doi:10.1017/s002211206800176x
15. Mysak L.A., Johnson E.R., Hsiem W.W. Baroclinic and barotropic instabilities of coastal currents // *Journal of Physical Oceanography*. 1981. Vol. 11, N 2. P. 209–230. doi:10.1175/1520-0485(1981)011<020>
16. Buchwald V.T., Adams J.K. The propagation of continental shelf waves // *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*. 1968. Vol. 305, N 1481. P. 235–250. doi:10.1098/rspa.1968.0115
17. Пелиновский Е.Н. *Гидродинамика волн цунами*. Нижний Новгород: ИПФ РАН, 1996. 156 с.
18. Didenkulova I., Pelinovsky E. On shallow water rogue wave formation in strongly inhomogeneous channels // *Journal of Physics A: Mathematical and Theoretical*. 2016. Vol. 49, N 19. P. 194001. doi:10.1088/1751-8113/49/19/194001
19. Ermakov A.M., Stepanyants Y.A. Transformation of long surface and tsunami-like waves in the ocean with a variable bathymetry // *Pure and Applied Geophysics*. 2019. Vol. 177, N 3. P. 1675–1693. doi:10.1007/s00024-019-02259-4
20. Rabinovich A.B., Thomson R.E. Evidence of diurnal shelf waves in satellite-tracked drifter trajectories off the Kuril Islands // *Journal of Physical Oceanography*. 2001. Vol. 31. P. 2650–2668. doi:10.1175/1520-0485(2001)031<2650:EODSWI>2.0.CO;2
21. Rabinovich A.B., Thomson R.E., Bograd S.J. Drifter observations of anticyclonic eddies near Bussol' Strait, the Kuril Islands // *Journal of Oceanography*. 2002. Vol. 58. P. 661–671. doi:10.1023/A:1022890222516
22. Булатов Н.В., Лобанов В.Г. Исследование мезомасштабных вихрей восточнее Курильских островов по данным метеорологических спутников Земли // *Исследование Земли из космоса*. 1983. № 3. С. 40–47.
23. Rogachev K.A. Rapid thermohaline transition in the Pacific western subarctic and Oyashio fresh core eddies // *Journal of Geophysical Research: Oceans*. 2000. Vol. 105, Iss. C4. P. 8513–8526. doi:10.1029/1999JC900330

24. *Kaneko H., Itoh S., Kouketsu S., Okunishi T., Hosoda S., Suga T.* Evolution and modulation of a poleward propagating anticyclonic eddy along the Japan and Kuril-Kamchatka trenches // *Journal of Geophysical Research: Oceans*. 2015. Vol. 120. P. 4418–4440. doi:10.1002/2014JC010693
25. *Prants S.V., Lobanov V.B., Budyansky M.V., Uleysky M. Yu.* Lagrangian analysis of formation, structure, evolution and splitting of anticyclonic Kuril eddies // *Deep Sea Research Part I: Oceanographic Research Papers*. 2016. Vol. 109. P. 61–75. doi:10.1016/j.dsr.2016.01.003
26. *Ефимов В.В., Рабинович А.Б.* О резонансных приливных течениях и их связи с континентальными шельфовыми волнами в северо-западной части Тихого океана // *Известия АН СССР. Физика атмосферы и океана*. 1980. Т. 16, № 10. С. 1091–1101.
27. *Пранц С.В.* Вихри глубоководных желобов северо-западной части Тихого океана: обзор // *Известия РАН. Физика атмосферы и океана*. 2021. Т. 57, № 4. С. 387–400. doi:10.31857/S0002351521040106

#### About the authors

**ГНЕВЫШЕВ Владимир Григорьевич**, РИНЦ Author ID: 298530, ORCID ID: 0000-0001-6654-5570, Scopus Author ID: AAZ-6352-2021, WoS ResearcherID: 6507346231, avi9783608@gmail.com

**ТРАВКИН Владимир Станиславович**, РИНЦ Author ID: 1023273, ORCID ID: 0000-0002-7254-9313, Scopus Author ID: 57509420800, WoS ResearcherID: HPE-4729-2023, v.travkin@spbu.ru

**БЕЛОНЕНКО Татьяна Васильевна**, РИНЦ Author ID: 66026, ORCID ID: 0000-0003-4608-7781, Scopus Author ID: 6507005889, WoS ResearcherID: K-2162-2013, btvlisab@yandex.ru